

FAST EDGE CORRECTED MEASUREMENT OF THE TWO-POINT CORRELATION FUNCTION AND THE POWER SPECTRUM

ISTVÁN SZAPUDI¹, JUN PAN^{1,2}, SIMON PRUNET³, AND TAMÁS BUDAVÁRI⁴*Draft version February 2, 2008*

ABSTRACT

We present two related techniques to measure the two-point correlation function and the power spectrum with edge correction in any spatial dimensions. The underlying algorithm uses fast Fourier transforms for calculating the two-point function with an heuristically weighted edge corrected estimator. Once the correlation function is measured, we estimate the power spectrum via numerical integration of the Hankel transform connecting the two. We introduce an efficient numerical technique based on Gauss-Bessel-quadrature and double exponential transformation. This, combined with our, or any similar, two-point function estimator leads to a novel edge corrected estimator for power spectra. The pair of techniques presented are the Euclidean analogs of those developed and widely used in cosmic microwave background research for spherical maps.

Subject headings: large scale structure — cosmology: theory — methods: statistical

1. INTRODUCTION

The two-point correlation function and its Fourier transform, the power spectrum, are the most measured statistics in large scale structure studies. Correlation functions can be measured naively with an $O(N^2)$ technique through cycling through pairs of objects. The advantage of that is that the edge corrected estimator of Landy & Szalay (1993) or Szapudi & Szalay (1998, 2000) (c.f. Equation 1) can be used for nearly optimal measurements. The naive counting of pairs can be sped up, if one is interested in measuring the correlation function on small scales, and reasonably large bins. In that case the double-tree algorithm by Moore et al. (2001) scales approximately as $O(N \log N)$. However, if all scales are considered, with high resolution, the tree-based algorithm has to split all nodes of the tree, thus it will degenerate into the $O(N^2)$ scaling of the naive algorithm.

Measurement of the power spectrum can be obtained directly with fast Fourier transforms (FFTs). The scaling of this algorithm $O(N \log N)$, where N corresponds to the dynamic range of the measurement. A particular advantage of the FFT is that the speed is independent of the maximum scale (or smallest wavenumber): scales up to the fundamental mode can be measured together with scales corresponding to the grid size. The resulting estimates, however, are convolved with the survey window. In cosmic microwave background (CMB) studies, such power spectra are termed pseudo-power spectra, and have been entirely replaced by edge corrected (or true) power spectra (e.g., Szapudi et al. 2001; Hivon et al. 2002)). In this paper we ask the question: can we take advantage of the FFT algorithms, and calculate the correlation function in $O(N \log N)$ time even on large scales? Can we exploit the edge corrected estimator

for the two-point correlation function, and estimate true (edge-corrected) power spectra? As we show next, the answer is affirmative to both questions, in complete analogy to results already obtained in CMB research.

In the next §2 we describe a new technique to measure edge corrected correlation functions using an FFT based algorithm, and a numerical technique to obtain edge corrected power spectra. We present numerical results in §3, and summarize and discuss them in the last §4.

2. DESCRIPTION OF THE TECHNIQUE

It is well known that the power spectrum is the Fourier transform of the two-point correlation function (Wiener-Khinchin theorem). In cosmic microwave background research, this basic idea was exploited by Szapudi et al. (2001) to obtain a fast method for estimating the angular power spectrum C_ℓ 's with fast harmonic transforms. A crucial intermediate step in their method was to estimate correlation functions (Szapudi et al. 2001). Although the window function can be inverted in harmonic space (Hivon et al. 2002), edge effect correction can be trivially obtained from the estimator Szapudi & Szalay (1998, 2000). This correspond to inverting the window matrix in pixel space, where it is diagonal. Our plan is to adapt this method for Euclidean space, arbitrary dimensions. The resulting technique is naturally divided into two steps: i) fast calculation of the two-point correlation function via FFT ii) calculation of the power spectrum via a numerical Hankel transform.

2.1. Two-point correlation function

The minimum variance estimator of Szapudi & Szalay (1998) is most natural when defined on a grid. If δ_i is the density field on grid point i , we can define the estimator as

$$\tilde{\xi}_\Delta = \sum_{i,j} f_{ij} \delta_i \delta_j / \sum_{i,j} f_{ij}, \quad (1)$$

where f_{ij} is the pair weight, and we assumed that $\langle \delta_i \rangle = 0$. Naive estimation with general pair weight is $O(N^2)$, where N is the number of pixels. As we will see,

¹ Institute for Astronomy, University of Hawaii, 2680 Woodlawn Dr, Honolulu, HI 96822, USA

² The School of Physics and Astronomy, University of Nottingham, University Park, Nottingham, NG7 2RD, England

³ Institut d'Astrophysique de Paris, CNRS, 98bis bd Arago, F-75014 Paris, France

⁴ Dept. of Physics and Astronomy, The Johns Hopkins University 3400 North Charles Street, Baltimore, MD 21218-2686

fast estimation is possible in the special case of the pair weight, where $f_{ij} = f_i f_j \delta_{i-j, \Delta}^{Kr}$, i.e. the weight is multiplicative, and depends only on a shift Δ . Here δ_{kl}^{Kr} is the Kronecker- δ , taking values of 1 when $k = l$, and 0 otherwise. The simplest example is flat weighting $f_i = 1$.

The correlation function without the normalization is a “raw-correlation” function of unnormalized counts. The normalization for an arbitrary complex geometry can be obtained by calculating the raw correlation function of the geometry, putting 1’s into valid grid points, and 0’s everywhere else. While our notation suggests $D = 1$ dimensions, in fact arbitrary dimension is included, simply by replacing indices, say i , with tuples of indices $(i_1, \dots, i_D)$.

Computationally, one needs to calculate pair summations of the form $\sum a_i a_{i+\Delta}$. These be reformulated to make use of Fast Fourier Transforms (FFTs), one of the fastest algorithms in existence. If $P(x) = a_0 + a_1 x + \dots + a_{n-1} x^{n-1}$ is a polynomial, and $\epsilon = e^{2\pi i/r}$ unit roots, the coefficients of a discrete Fourier transform of the series a_i can be defined as

$$\hat{a}_k = P(\epsilon^k) = a_0 + a_1 \epsilon^k + \dots + a_{n-1} \epsilon^{(n-1)k}. \quad (2)$$

Direct calculation confirms that $\sum a_k b_{k+\Delta}$ can be calculated by Fourier transforming the series a_i and b_i , multiplying the resulting $\hat{a}_k \hat{b}_k^*$, and finally inverse Fourier transforming back. This simple observation is the discrete analog of the Wiener-Khinchin theorem. The Fourier space quantity arising in the above algorithm is the discrete, Euclidean “pseudo-power spectrum”. Note that in fact all previous direct measurements of the power spectrum using Feldman et al. (1994) type estimators measured pseudo power spectrum. We will see in the next subsection, how to obtain the (edge corrected) power spectrum instead from the correlation function.

Using the above observation, we can put together a fast algorithm to calculate two-point correlation function on in D dimensions with the following steps: i) placing the points in a sufficiently fine grid, storing the value $N_{\mathbf{k}}$, the number of objects (e.g. galaxies) at (vector) grid point $\mathbf{k}$, (this step is omitted if the density field is a given, such as Euclidean approximation of CMB maps, etc.) ii) calculating fluctuations of the field by $\delta = (N - \langle N \rangle) / \langle N \rangle$ for each grid point iii) possible weighting each point with a minimum variance weight (e.g. J_3 weighting, or Feldman et al. (1994) weighting), iv) discrete Fourier transform with a fast FFT engine, v) multiplying the coefficients vi) Fourier transform back. The same procedure is followed for the mask with zero padding large enough (i.e. larger than the largest bin of the correlation function) to avoid aliasing effects. Finally, the raw correlation function is divided with the correlation function of the mask, according to Equation 1.

The result will be an inhomogeneous correlation function $\tilde{\xi}(\mathbf{r})$, depending on vector shifts $\mathbf{r}$. To obtain the traditional correlation function one has sum the result in rings/spheres for $D = 2, 3$, respectively.

2.2. Power Spectrum

The power spectrum is a D -dimensional Fourier transform of the two-point correlation function. Using rotational invariance, this will become a Hankel transform in $D = 2, 3$ dimensions:

$$\xi(r) = \int \frac{k dk}{2\pi} P(k) J_0(kr) \quad 2D,$$

$$\xi(r) = \int \frac{k^2 dk}{2\pi^2} P(k) j_0(kr) \quad 3D, \quad (3)$$

where J_0 and j_0 are ordinary and spherical Bessel functions, respectively. As well known the Hankel transform is self inverse, and our plan is to use the inverse of the above formulae to obtain an estimator of the power spectrum.

In practice, it is a delicate numerical procedure to calculate a Hankel transform numerically. We use the quadrature formulae of Ogata (2005), which uses both the roots of Bessel functions, as well as the double exponential transformation of Ooura & Mori (1999). Since these techniques are fairly new developments in numerical mathematics, we summarize the recipe here. Explicitly, to integrate over an arbitrary function f multiplied with a Bessel function we use the formula

$$\int_0^\infty f(x) J_\nu(x) \simeq \pi \sum_{k=1}^\infty w_{\nu k} f(\pi \psi(hr_{\nu k})/h) \times J_\nu(\pi \psi(hr_{\nu k})/h) \psi'(hr_{\nu k}), \quad (4)$$

where $r_{\nu k}$ are the roots of the Bessel function J_ν , $\psi(t) = t \tanh(\frac{\pi}{2} \sinh t)$ the double exponential transformation, h is the step of the integration (similarly to that of the trapezoidal rule). The weights are calculated as $w_{\nu k} = 2/(\pi^2 r_{\nu k} J_{\nu+1}(\pi r_{\nu k}))$. For the inverting the two-point function $\nu = 0$ and $\nu = 1/2$ Bessel functions should be used for 2 and 3 spatial dimensions according to Eqs 3.

3. RESULTS




FIG. 1.— The two-point correlation function measured with the Moore et al. (2001) tree-based algorithm (black symbols with errorbars) vs. our Fourier based approach (red/gray) in a set of VLS simulations (see description in text). The agreement is virtually perfect, despite the difference in the methods and binning. For comparison, we also show a separate measurement of the two-point function where (erroneously) periodic boundary conditions were assumed (solid line). This can be considered as a measurement without proper edge correction.

The above algorithm to measure the correlation function was implemented in C using the FFTW3 package for two and three dimensions. The two-dimensional version of the algorithm was extensively tested in Budavári et al. (2003), the first application to measure the angular correlation function in the SDSS. In what follows, we focus on three-dimensions.

To test our algorithm for the two-point correlation function, and the power spectrum, we performed measurements in Λ CDM simulations by the Virgo Supercomputing Consortium Jenkins et al. (1998). We used outputs of VLS (Very Large Simulation) with the following cosmological parameters: $\Omega_m = 0.3$, $\Omega_v = 0.7$, $\Gamma = 0.21$, $h = 0.7$ and $\sigma_8 = 0.9$. In order to estimate measurement errors, we divided the VLS simulation into eight independent subsets, thus each subset was $(239.5h^{-1}\text{Mpc})^3$ cube. Note that thus sub-cubes are *not* subject to periodic boundary conditions, thus, although their geometry is simpler than a typical galaxy survey, they are subject to edge effects, unlike a full simulation cube.

We have measured the correlation function in logarithmic bins in the range of $0.3 - 162h^{-1}\text{Mpc}$ with our implementation of the algorithm **eSpICE** (Euclidean version of SpICE, the spatially inhomogeneous correlation estimator, Szapudi et al. (2001)) Measurement of the correlation function on a 768^3 grid takes about 15 minutes on one Opteron processor. In addition, we also performed control measurements with the Moore et al. (2001) tree-based algorithm in logarithmic bins from $0.1 - 38h^{-1}\text{Mpc}$ scales. The required CPU time of this latter algorithm increases drastically on large scales, hence we stopped at smaller scales. Even though the binning of the two runs were not identical, the two algorithms produced virtually identical curves as shown on Figure 1. In addition, we have measured the correlation function assuming –erroneously, according to our previous discussion– periodic boundary conditions. The difference between the results based on periodic and non-periodic boundary conditions demonstrate that edge effects are already significant on $40h^{-1}\text{Mpc}$ scales in a $239.5h^{-1}\text{Mpc}$ size simulation.

The next step was calculating the edge corrected power spectrum. First, we have performed a cubic spline interpolation on the measured correlation function, to turn the discrete points into a function, which can be sampled at the roots of Bessel functions. Then we applied the recipe of Equation 4 for $\nu = 1/2$ to integrate the first of Eqs. 3. We used $h = 1/32$ and truncated the sum at $N = 200$; doubling these values made no significant effect on our results. We have checked the accuracy of our integration routine with $N = 10000$ point integration according to the direct double exponential integration by Oura & Mori (1999), and with **Mathematica**. This showed that this fast 200 point integration is better than 0.1% accurate, which was good enough for our purposes. Note that with some care, **FFTLog** (Hamilton 2000) could probably be adapted to the same purpose, although the aliasing and numerical effects are more delicate.

The results are shown on Figure 2. We have applied the inversion both to the results from **eSpICE**, and from the tree based algorithm. We used a scale range conservatively as $(k_{\min}, k_{\max}) = 2\pi/r_{\max}, 1/r_{\min}$. In addition we plotted the linear theory and non-linear Smith et al. (2003) power spectra. The agreement of our reconstruction with the phenomenology is nearly perfect.

4. SUMMARY AND DISCUSSIONS

We have presented a fast Fourier based algorithm, which is complementary to the tree-based algorithm of Moore et al. (2001). While the latter is ideal for mea-




FIG. 2.— The power spectrum extracted from the two-point correlation function using the method described in the text. The green and red lines show the linear and non-linear Smith et al. (2003) power spectra. The other symbols and lines correspond to Figure 1.

suring the correlation function on small scales, taking advantage of a KD-tree construction, our grid based approach is ideal for large scales. Our algorithm has the advantage that it does not slow down on large scales, although the grid has a smoothing effect on the correlation function on scales a few times the grid spacing.

An interesting side effect of our algorithm is that as an intermediate step it produces inhomogeneous correlation functions of vector shifts $\xi(\mathbf{r})$. These have been used in Budavári et al. (2003) to filter out systematic effects from the drift scanning operations of the camera. In three dimensions, this could be used for estimation of the redshift space correlation function in the distant observer approximation, for narrow, deep redshift surveys.

We also presented a method to turn a measured correlation function into an edge corrected power spectrum. This is, apart from numerical details, a direct Euclidean generalization of the spherical method by Szapudi et al. (2001). This new technique is independent on the details of how the two-point function is estimated: our Fourier based approach, a tree-based algorithm, or even a direct N^2 algorithm can be used to measure the two-point function. The inversion for the power spectrum should work just the same. In the present work we did not correct for grid effects in the inverted power spectrum, but it should scale as $W(k)^2 = \text{sinc}(k)^{2D}$ in D dimensions.

Were we measuring the correlation function in linear scales (a built in option for **eSpICE** which does not influence the speed), we would have obtained $P(k)$ in linear bins. The maximum $r_{\max}$ at which the correlation function is measured, determines the resolution, and the smallest $k_{\min} \simeq \Delta k \simeq 1/r_{\max}$ for $P(k)$, while the resolution and the smallest $r_{\min}$ at which the correlation function is measured, determine $k_{\max} \simeq 1/r_{\min}$.

In accordance with the overwhelming majority of practical measurements in the past, we choose to measure the correlation function in logarithmic bins. As a result

we obtained $P(k)$ in approximately logarithmic bins in k . To see this, one has to note that a fixed resolution corresponding to a particular k in the power spectrum, corresponds to a maximum r at which the correlation function is measured at this resolution. This maximum translates into an effective resolution Δk at this particular k . This qualitative argument can be made more rigorous with calculation of the effective window function if needed.

Large scale structure studies, up until now, have estimated “pseudo power spectra” (a term borrowed from cosmic microwave background research) i.e. one convolved with the survey geometry. The potential of detecting baryonic oscillations in the measurement of the power spectrum motivates the need for measuring power spectra with the highest resolution possible. In particular, if one of the survey dimensions is much smaller than the other two, the effective convolution kernel will be much wider than desirable. In the above we demonstrated that the power spectrum can be recovered from steps not any more complicated than the measurement of the corresponding pseudo power spectrum. While the information content cannot be improved with the inversion (e.g. Efstathiou 2004), it is immensely useful when comparing measurements from different surveys, from different geometries, as it has been common in CMB research.

Note that the present technique can be used to esti-

mate C_ℓ ’s in the flat sky approximation, using the asymptotic behavior of the Legendre polynomials $P_\ell(\cos\theta) \simeq \sqrt{\frac{2}{\pi\ell\sin\theta}} \cos((\ell+1)\theta - \pi/4)$, and the Bessel functions $J_0(z) \simeq \sqrt{\frac{2}{\pi z}} \cos(z - \pi/4)$. The correlation function can be measured with **eSpICE**, and integration of Equation 3 gives the C_ℓ ’s with approximately $k \simeq \ell + 1/2$. Generalization of the proposed computational and inversion techniques to gravitational lensing follows directly from the generalization **SpICE** to CMB polarization, such as Chon et al. (2004). Finally, generalization of Equation 3 for a relation between the three-point function and the bispectrum (Szapudi 2004) together with a fast algorithm to measure the three-point function yields a new, edge corrected method to measure the bispectrum. These generalizations will be published elsewhere.

Our **eSpICE** implementation will be made public upon acceptance of this paper.

I thank Alex Szalay for stimulating discussions, Josh Hoblitt and Gang Chen for computing help. IS and JP was supported by NASA through AISR NAG5-11996, and ATP NASA NAG5-12101 as well as by NSF grants AST02-06243, AST-0434413 and ITR 1120201-128440. JP acknowledges support of PPARC Grant number PPA/G/S/2000/00057.

REFERENCES

- Budavári, T., et al. 2003, *ApJ*, 595, 59
 Chon, G., Challinor, A., Prunet, S., Hivon, E., & Szapudi, I. 2004, *MNRAS*, 350, 914
 Efstathiou, G. 2004, *MNRAS*, 349, 603
 Feldman, H. A., Kaiser, N., & Peacock, J. A. 1994, *ApJ*, 426, 23
 Hamilton, A. J. S. 2000, *MNRAS*, 312, 257
 Hivon, E., Górski, K. M., Netterfield, C. B., Crill, B. P., Prunet, S., & Hansen, F. 2002, *ApJ*, 567, 2
 Jenkins, A., et al. 1998, *ApJ*, 499, 20
 Landy, S. D., & Szalay, A. S. 1993, *ApJ*, 412, 64
 Moore, A. W., et al. 2001, in *Mining the Sky*, 71
 Ogata, H. 2005, *Publ. RIMS, Kyoto Univ.*, Accepted
 Oura, T., & Mori, M. 1999, *J.Comp.Appl.Math*, 112, 229
 Smith, R. E., et al. 2003, *MNRAS*, 341, 1311
 Szapudi, I. 2004, *ApJ*, 605, L89
 Szapudi, I., Prunet, S., & Colombi, S. 2001, *ApJ*, 561, L11
 Szapudi, I., Prunet, S., Pogosyan, D., Szalay, A. S., & Bond, J. R. 2001, *ApJ*, 548, L115
 Szapudi, I., & Szalay, A. S. 2000, *Annales de l’I.S.U.P.*, 1
 Szapudi, S., & Szalay, A. S. 1998, *ApJ*, 494, L41